

:  
/ / :  
/ / :

± /  
/ ± / , / ± / , /  
( )  
( ) -

**B A** ( / ± / )  
( / ± / - / ± / )  
(P < / )



'

— .( , , , , , )

.( )

— —

— .( , )

'

)

/

.( , )(

.( )

.( , )

'



.( )

( )

.( )

.( )

( **B A** )

---

1- Cardiovascular Fitness Ratio



( )

( )


, ( **Seca** )

**(ACSM)**

.( )

, ( )

( )



C°                    %                    )                    (                    )                    .                    (                    )

-                    .                    -                    /                    ,                    ,                    -                    .                    ,                    ,

)( )  
: (SEE = % /    r = /  
CFR = / - / \*                    (                    )

(C                    )                    /

:  
CFR = / - / \*                    (                    )

(B                    )                    /  
-                    ,                    (A                    )



:

**A: CFR** = / - / \* ( )

**B: CFR** = / - / \* ( )

)

( ) (r<sup>2</sup> r,SEE)

( ) -

-

**MedCalc(8.2.1.0)**


( / )	( )	( )	( )
/ ± /	/ ± /	/ ± /	/ ± /

**A** ) , ( )




- ( **B**

---

1- Agreement  
2- Bland – Altman (1986)



			%		
/	/	/	/ /	/	-C
/	/	/	/ /	/	-A
/	/	/	/ /	/	-B



( ) A

( )

( ) (B)

(X)

( )


(y)



-



( )  
( ) ( % )



( )

-

-

( )  
( )

-

:


$$- = / - / * ( )$$

A

$$- = / - / * ( )$$

B

$$- = / - / * ( )$$



( )B

( )A

( )C

-

---

% , -  
( )

	%		
<b>P&lt; /</b>	/ /	/	<b>C</b>
<b>P&lt; /</b>	/ /	/	<b>A</b>
<b>P&lt; /</b>	/ /	/	<b>B</b>

**T** , , -  
( )

	<b>T</b>			
<b>P&lt; /</b>	/	/	/	<b>C</b>
<b>P&lt; /</b>	/	/	/	<b>A</b>
<b>P&lt; /</b>	/	/	/	<b>B</b>

-

- (C )  
( )

- % .( )

- (C )  
( )

(C )

-

\_\_\_\_\_

-  
)  
, ( ) , ( ) , ( ) , ( ) ( )  
( )

-  
( , , , , )  
( )  
, ( )

-  
(A )  
) % ( )  
, ( )

- ( ) (A )  
( ) (B )  
, ( ) (A )

(A ) / (B )  
( )  
(A )



( ) ,

-

( , , , , )

,A

( )

,

.

,A

,

A

.

,

-

-

-

( )

-

.

.

"( ) ,

, , , , ,

:( - )

- 
- 
3. American College of Sports Medicine. (2000). "ACSM'S guidelines for exercise testing and prescription (6<sup>th</sup> ed)". Philadelphia PA: Lippincott Williams & Wilkins :91.
4. American College of Sport Medicine. (1990). "ACSM'S guidelines for exercise testing and prescripting. 4<sup>th</sup> edition, lea and febiger, P:148.
5. Arrais GA, Wigle DT, and Mao Y. (1992). "Risk assessment of physical activity and physical fitness in the Canadian Health survey follow up study". *J Clin Epidemiol* ; 45: PP:419-28.
6. Blair S.N. Kohl H.W. (1989). "Physical fitness and all cause mortality : a prospective study of Health men and women". *J of the American Medical Association*; 262 : PP:2395-2401.
7. Blair , S.N. Kohl, H.W., Gordon, N.F., & Paffenbarger, R.S. (1992). "How much physical activity is good for health? Annual Review of public Health, 13, PP:99-126.
8. Blair S.N., Ming, W., and chong, D.L (1998). "Cardiorespiratory fitness determined by exercise heart rate as a predictor of mortality in the aerobic center longitudinal study". *J Sport sci*; 16, PP: S47-S55.
9. Bland, M.J. & D.G. Altman, (1986). "Statistical methods for assessing agreement between two methods of clinical measurement". *The Lancet*, PP:1307-310.
10. Bradshaw DI, George JD, Hyde A, Lamonte MJ, Vehrs PR, Hager RL, Yanowitz FG. (2005). "An accurate VO<sub>2</sub>max nonexercise regression model for 18-65-year-old adults". *Res Q Exerc Sportt.*;76(4) : PP:426-32.
11. Byrne NM. Hills AP, Hunter GR, Weinsier RL, Schutz Y. (2002). "The metabolic equivalent : One size does not fit all". *J Appl Physiol*. 14; [Epub ahead of print].
12. Donald C. Dibloo and Jeffery K. Moffit (2003). "A comparison of bioelectrical impedance and NEAR-Infrared Interact ance to skinfold measures in determining minimum wrestling weight in collegiate wrestlers". *International Electronic Journal Vol. 6. No.2* : PP:26-36.
13. Ekelund L.G., Haskell WL. (1988). "Physical fitness as a predictor of cardiovascular mortality in asymptomatic North American men". *N Eng J Med*, 319 : PP:1379-84.

- 
14. Franz X. Kleber . (2004). "The predictive value of cardiorespiratory fitness". *European Heart Journal* 25, PP: 1374-1375.
15. Gibbons L, Blair SN, Kohl HW, Cooper K (1989). "The safety of maximal exercise testing". *Circulation*; 80 : PP:846-852.
16. Hien H.O. (1992). "Physical fitness or physical activity as a predictor of ischemic heart disease". *J intern Med*, 232; PP:471-9.
17. James D.G., Pat R.V. & Garth J.B. (2000). "A modified submaximal cycle ergometer test designed to predict treadmill  $VO_{2max}$  , Vol, 4, PP:229-243.
18. Lie H. Mundal R.and Erikssen J. (1985). "Coronary risk factor and incidence of coronary death in relation to physical fitness". *Eur heart J*, 6 : PP:147-57.
19. Seefeldt V, RM Malina, and MA Clark. (2002). "Factors affecting levels of physical activity in adults". *Sports Med*. 32; PP:143-168.
20. Slattery, M.L. Jacobs, D.R.Jr. (1988). "Physical fitness and cardiovascular disease mortality"; the US railroad study *Am J Epidemiol*, 127;PP:601-10.
21. Spangler, R.D. (1970). "A submaximal exercise electrocardiographic test as method of determining occult ischemic heart disease". *Am Heart J* 80: PP:725-58.
22. Thomas, J.R.& J.K. Nelson . (1990). "Research methods in physical activity". 2th ed, Chmpaign : *Human Kinetics*. PP:129-178.
23. Wei M.JB Kamper T, CE Barlow, MZ Nichaman, LW Gibbons, RS Paffenbarger Jr, ans SN Blair. (1999). "Relationship between low cardiorespiratory fitness and mortality in normal-weight, overweight, and obese men". *JAM A*. 282; PP:1547-1553.
24. Young DZ, Lampert S, Graboys TB, Lown B. (1984). "The safety of maximal exercise testing in patients at high risk for ventricular arrhythmia; 70;PP:184-191.